

MPW-485265

by Academic 2025

General metrics

67,692

characters

9,578

words

524

sentences

38 min 18 sec

reading
time

1 hr 13 min

speaking
time

Score



This text scores better than 87%
of all texts checked by Grammarly

Writing Issues

209

Issues left

27

Critical

182

Advanced













Plagiarism

This text hasn't been checked for plagiarism

Writing Issues

34

Correctness

2	Improper formatting	
6	Misspelled words	
4	Text inconsistencies	
1	Citation style options	
5	Incorrect punctuation	
2	Incorrect verb forms	
2	Incorrect noun number	
5	Determiner use (a/an/the/this, etc.)	
2	Conjunction use	
2	Closing punctuation	
1	Wrong or missing prepositions	
2	Ungrammatical sentence	

113

Style guide

113	Custom suggestions with replacements	
-----	--------------------------------------	---

62

Clarity

7	Intricate text	
52	Paragraph can be improved	
2	Wordy sentences	
1	Passive voice misuse	

Unique Words

Measures vocabulary diversity by calculating the percentage of words used only once in your document

18%

unique words

Rare Words

Measures depth of vocabulary by identifying words that are not among the 5,000 most common English words.

46%

rare words

Word Length

Measures average word length

5.6

characters per word

Sentence Length

Measures average sentence length

18.3

words per sentence

MPW-485265

Relationship Between Childhood Obesity and Anxiety Among U.S. Children: The Moderating Effects of Gender and Urbanicity

Doctor Of Public Health

By

Mph¹, Capella University, 2018

BS, National Graduate School, 2015

Doctoral Study Submitted in Partial Fulfillment
of the Requirements for the Degree of
Doctor of Public Health

Walden University

06/30/ 2026

Abstract

The prevalence of childhood obesity² (2-19 years) in the United States (U.S.) is approximately 20%, and it is³ associated with physical and mental health complications. Although⁴ obesity⁵ is associated with mental health problems, there is limited research on the moderating effects of gender and urbanicity on the association between obesity⁶ and anxiety. This⁴ quantitative, cross-sectional, secondary data analysis of the 2022-2023 National Survey of Children's Health examines this link among children aged 10-17. The⁴ research engages with the Social Ecological Model to examine the interaction of gender and urbanicity on the association between body mass index (BMI) classification and parent-reported anxiety, adjusting for depression, age, race/ethnicity, and socioeconomic⁷ status. Weighted⁴ logistic regression models with interaction are used. Results⁴ will guide culturally tailored interventions for obesity⁸ and anxiety among at-risk groups, such as rural areas and gender-specific groups.

Keywords: childhood obesity⁹, anxiety, National Survey of Children's Health, gender, urbanicity, Social Ecological Model, logistic regression, moderation

Dedication

This¹⁰ is an optional page for a dedication. If⁴ you include a dedication, use regular paragraph spacing as shown here (not centered, italicized, or otherwise formatted). If⁴ you do not wish to include this page, delete the heading and the body text.

Acknowledgments

¹¹
----- This is an optional page for acknowledgments. ⁴
----- It is a nice place to thank the faculty, family members, and friends who have helped you reach this point in your academic career.

No page number appears on any of the pages up to this point. ⁴
----- If you do not wish to include this page, delete the heading and the body text, but leave the section break that you see below this text.

iii

Table of Contents

List of Tables iii

List of Figures iv

Section 1: Foundation of the Study and Literature Review (APA Level 0 Heading)

1

Topic of the Study 1

Background 2

Problem Statement 5

Theoretical Framework 14

Definitions 19

Assumptions 21

Scope and Delimitations 22

Limitations 23

Section 2: Research Design and Data Collection 27

Introduction 27

Research Design and Rationale 27

Methodology 28

Section 3: Presentation of the Results and Findings 37

Section 4: Application to Professional Practice and Implications for

SocialChange¹² 38

References 39

Appendix A: Title of Appendix 42

List of Tables

Table 1. A⁴ Sample Table Showing Correct Formatting 5

Table 2. Another⁴ Sample Table 5

List of Figures

Figure 1. Figure⁴ Title 6

Figure 2. Another⁴ Figure Title 6

13

Section 1: Foundation of the Study and Literature Review (APA Level 0 Heading)

Introduction

The problem of childhood obesity¹³ is one of the most acute issues in the 21st-century United States public health. Data⁴ provided by the Centers for Disease Control and Prevention (CDC) indicate that obesity¹⁴ rates among children and

adolescents aged 2-19 reflected a worrisome increase because in 2017-2020, it was estimated to be about 20% compared to 16.9% in 2009-2010 (Contreras et al., 2021; Crouch et al., 2023). In addition to the documented physical outcomes, such as higher risks of type 2 diabetes, cardiovascular disease, and musculoskeletal disorders, childhood obesity is now increasingly viewed as an antecedent and predictor of abnormal mental health conditions (especially anxiety and depression). The physical and mental health of children are closely interconnected, requiring strict scientific consideration, and mental disorders during childhood have long-term implications during adulthood.

Although there is a growing body of literature on mental health issues in youth related to obesity, there are still some gaps in the knowledge about the influence of demographic and geographic factors on such relationships. In particular, there are not many studies that have investigated the moderating effects of gender and urbanicity concomitantly, which are two factors with different, and possibly interactive effects on the obesity-anxiety relationship. The knowledge of the rural and urban residence or gender identity as either reinforcing or reducing the psychiatric burden of childhood obesity is critical to the formulation of specific, context-specific interventions. This paper fills that gap by using a nationally representative sample, the 2022-2023 National Survey of Children's Health (NSCH), to test the joint moderation effect between gender and urbanicity on the relationship between children's obesity and anxiety in the United States aged between 10 and 17.

In this section, the theoretical and empirical background of the study is presented. It features following key subsections: (a) a review of the background literature that determines the scope and importance of childhood obesity and the mental health correlates of obesity; (b) a clear statement of the research problem and the research gap identified; (c) the purpose of the study; (d) the

research questions and hypotheses; (e) the theoretical framework used to conduct the research; and (f) the nature of the study such as the research design, sources of data, variables, and method of analysis. The given structure is created to build the logical flow between a sociologically stated problem and a scientifically based research question.

Background

Childhood obesity²³ is a multifactorial disorder, with a complex association of biological, behavioral, and environmental variables. It⁴ has grown significantly in the United States over the last 40 years due to changes in dietary choices, a decline in physical activities, screen time, and wider social determinants of health encompassing food insecurity, poverty, and restricted availability of safe recreational areas (Contreras et al., 2021; Forster et al., 2023).^{24 15 4} Rural populations are especially underprivileged (with less access to healthy food choices, fewer recreational opportunities, and greater socio-economic⁷ vulnerability) (Crouch et al., 2023; Foster et al., 2020).^{15 4} Studies based on NSCH data have observed a steadily higher risk of rural children being overweight or obese than children in urban areas, irrespective of demographic factors such as race/ethnicity and household income (Crouch et al., 2023).¹⁵

Childhood obesity²⁵ has a significant mental health burden that is multidimensional in nature. Obese⁴ children are at a higher risk of anxiety, depression, low self-esteem, social isolation, and peer victimization, which worsen their weight-related behaviors and introduce a vicious cycle of physical and psychological suffering (Beltran-Garrayo et al., 2023; Galler et al., 2024; Kokka et al., 2023).^{15 4} As Beltran-Garrayo et al. (2023) showed, obese youths displayed much greater scores of anxiety mediated by low body esteem and high levels of teasing, indicating that psychosocial mechanisms of excessive weight and mental health morbidity interrelationship exist. These⁴ dynamics

were further explained in a longitudinal study by Beltran-Garrayo et al. (2024), which revealed that the relationship between childhood obesity²⁶ and depressive symptoms in adolescents was mediated by body esteem and that even after adjusting for gender, the relationship is still strong.

Psychiatric comorbidity of obesity²⁷ in pediatric patients cuts across several diagnostic categories. A⁴ large clinical cohort study of children, adolescents, and young adults with obesity²⁸ and comorbid mental health disorders, Galler et al. (2024) discovered that anxiety disorders were one of the most widespread psychiatric comorbidities in the population. Adolescents⁴ with comorbid mental health also exhibited worse treatment outcomes of lifestyle-based weight management interventions; in this case, psychiatric burden is directly associated with intervention efficacy. Population-level⁴ systematic reviews support these clinical findings. Kokka⁴ et al. (2023) reviewed 14 cross-sectional studies, of which 9 reported statistically significant associations between psychiatric disorders and obesity²⁹. The⁴ most consistent evidence emerged for affective disorders, particularly depression, though the association was meaningful but weak. Anxiety⁴ disorders showed the strongest — albeit still weak — support across the reviewed studies.

The cross-cultural consistency of these associations has also been confirmed by epidemiological research. A⁴ systematic review of childhood obesity³⁰ and mental health outcomes by Ranjani et al. (2021) found that the connection³¹ between obesity³² and mental disorders, namely anxiety and depression, occurs repeatedly in the diversity of different cultural backgrounds, which confirms the validity of these relationships in not only particular nation-specific but also universal settings. A⁴ recent cross-sectional study conducted in Europe by Papadimitriou et al. (2025) demonstrated strong evidence (nearly twice as many depressive and anxiety symptoms) of overweight and obese children aged

6-9 years compared to normal-weight children, and that the strength of those associations varied according to gender and urban versus rural dwellers. The⁴ gender and urbanicity effects of moderation observed in the European sample indicate that this can be applied to the U.S. pediatric population, but this has not been rigorously assessed using nationally representative U.S. data.

This fact supports the role of gender as a potential moderator of the relationship between obesity³³ and mental health: it is believed that boys and girls might respond to obesity-related stigma and body dissatisfaction more or less (Beltran-Garrayo et al., 2024; Forster et al., 2023).^{24 15 4} In a large German cohort (aged 4-18 years), Forster et al. (2023)²⁴ discovered³⁴ that the relationships between BMI and mental health outcomes, such as behavioral problems and health-related quality of life, depended on both gender and socioeconomic⁷ status, making it important that³⁴ these moderators be considered together.³⁴ Likewise⁴, parental mental health has been identified as a significant contextual variable, differentially predicting obesity³⁵ risk among children across rural and urban environments and income groups (Foster et al., 2020),¹⁵ underscoring the relevance³⁷ of considering family and environmental context in obesity studies in³⁶ children.³

Regardless of³⁸ this accumulating evidence, there is³⁸ an identifiable gap in the literature. The⁴ majority of the available research focuses on the occurrence of obesity³⁹ and anxiety or depression without considering geographic and gender-based moderation, and only a handful of research studies use nationally representative U.S. data to simultaneously test the hypothesis of these moderators. The⁴ NSCH offers a distinct and strong platform toward such analysis since it has validated parent-reported indicators of child health status, mental health diagnosis, BMI-based obesity⁴⁰ status and some demographic characteristics such as gender and urbanicity. This⁴ research will

address this gap and extend the evidence base for targeted public health interventions.

Problem Statement

The research issue that the study seeks to focus on is that there is no detailed population-level knowledge on the relationship between anxiety and childhood obesity⁴¹ in the U.S. among children aged between 10 and 17, and the extent to which gender and urbanicity moderate this relationship. The⁴ prevalence of childhood obesity⁴² has risen to about 20% in 2017-2020 compared to about 16.9% in 2009-2010 (Contreras et al., 2021; Crouch et al., 2023)¹⁵; anxiety disorders are considered one of the most common psychiatric diseases in the children population, with the estimated rate of 7-10 per cent across the country. However⁴, the literature on the relationship between obesity⁴³ and anxiety has not studied the relationship in a systematic manner that incorporates both gender and urbanicity as factors. Thus⁴, there are critical gaps in the evidence base for intervention targeting.

The existing literature reports strong correlations between anxiety and obesity⁴⁴ across different study designs and geographic regions (Beltran-Garrayo et al., 2023; Galler et al., 2024; Kokka et al., 2023; Papadimitriou et al., 2025; Ranjani et al., 2021)¹⁵. The⁴ main limitations of these studies are mostly a non-representative sampling frame, single-moderator studies conducted in isolation, and an out-of-the-U.S. context. The⁴ NSCH data have been mostly⁴⁶ utilized⁴⁶ in studies on the prevalence of obesity⁴⁵ and rural-urban differences in weight status (Crouch et al., 2023; Foster et al., 2020)¹⁵, without analyzing mental health outcomes as core dependent variables and the moderate effects of gender and urbanicity on each other⁴⁶. The⁴ absence of evidence regarding combined moderation restricts the capacity of practitioners working in the sphere of public health, school health, and policymakers in creating gender-

sensitive and geographically focused interventions, which would be able to respond to the needs of particular risk groups of anxious obese children. This research is based on the current literature in the area of childhood obesity and mental health, specifically extending two streams of previous research: moderated mediation models (Beltran-Garrayo et al., 2024) and rural-urban comparative studies (Contreras et al., 2021; Crouch et al., 2023). It develops these frameworks in two important ways - by using a nationally representative U. S. sample and by introducing a dual moderation, that is, two variables simultaneously moderate the relationship under study. This is a logical progressive step in the field. Its results have practical implications on research studies on obesity and anxiety in children in the U.S., including the reinforcement of the evidence base, informing the healthcare practice, and assisting in preventing the co-occurrence of obesity and anxiety in children.

Purpose of the Study

This study is a quantitative, cross-sectional study that tests the relationship between childhood obesity and anxiety among children aged 10-17 years in the U.S., using secondary data from the 2022-2023 National Survey of Children's Health (NSCH). In particular, the research will focus on identifying whether the connection between childhood obesity (independent variable: BMICLASS, operationalized as BMI 95th percentile per CDC growth charts) and parent-reported anxiety diagnosis (dependent variable — that is, whether the strength or direction of this relationship differs by gender). The secondary objective is to test the relationships between childhood obesity and depression as a dependent variable using the same set of moderators and covariates. The field-based products of this study, such as a policy brief memo, a community health intervention plan, a visual illustration of the intervention

framework⁵⁹, and a focused fact sheet, will be directly important to the policymakers, school-based public health practitioners, pediatric health practitioners and community-based organizations that deal with the rural and urban children.⁴ The results of this study will allow allocating the resources available to the community in terms of obesity⁶⁰ and anxiety more accurately and efficiently, as well as will⁶¹ facilitate the creation of culturally and geographically specific intervention⁶², by clarifying the data about which subgroups of children (e.g., rural girls or urban boys) are at the highest risk of combined⁶³, obesity^{64 65}, and anxiety.

Research Question(s) and Hypotheses

This study is guided by two primary research questions, each accompanied by a null and alternative⁶⁶ hypothesis:

RQ1: What is the association between childhood obesity⁶⁷ and anxiety among U.S. children ages 10–17 when the moderating effects of gender and urbanicity are taken into account, controlling for depression, age, race/ethnicity, and socioeconomic⁷ status?

H₀₁: There is no statistically significant association between childhood obesity⁶⁸ and anxiety after controlling for covariates and including gender and urbanicity as moderators.

H₁₁: There is a statistically significant positive association between childhood obesity⁶⁹ and anxiety after controlling for covariates and including gender and urbanicity as moderators.

RQ2: What is the association between childhood obesity⁷⁰ and depression among U.S. children ages 10–17 when the moderating effects of gender and urbanicity are taken into account, controlling for anxiety, age, race/ethnicity, and socioeconomic⁷ status?

H₀₂: There is no statistically significant association between childhood obesity⁷¹ and depression after controlling for covariates and including gender and urbanicity as moderators.

H₁₂: There is a statistically significant positive association between childhood obesity⁷² and depression after controlling for covariates and including gender and urbanicity as moderators.

The following infographic summarizes the alignment between research questions, variables, data sources, and analytic procedures:

Figure 1 Research Design Alignment Summary

Theoretical and/or Conceptual Framework

Theoretical Framework

The theoretical basis of this research is the social-ecological model (SEM) of Bronfenbrenner (1979), which offers a multi-level⁷³ framework for explaining health and human development.⁷⁴ The SEM theorizes⁴ human development⁷⁵ and health outcome⁷⁷ are the results of a nested system of interacting environmental systems: microsystem (immediate personal environment), mesosystem (relationships among microsystems), exosystem (external systems whose influence influences the individual), macrosystem (cultural, policy and societal contexts),⁷⁸ and chronosystem (temporal aspect of environmental change).⁷⁶ These⁷⁸ systems within a system dynamically interact to influence health behaviors and lifelong outcomes.

The key theoretical propositions of the SEM applicable in this research are as follows: (a) personal characteristics do not alone determine the individual health outcomes but rather it is a complex interaction of various levels of environmental context that determines the variation in health outcomes; (b) the interplay of individual and community-level factors (such as gender versus

urbanicity) is a key determinant of differential health consequences; and (c) effective public health interventions have to simultaneously address factors at different levels of the ecological context to produce a sustained change. These propositions lead directly to the study's analytic approach, in which gender is treated as an individual-level moderator, urbanicity as a community-level moderator, and how the two interact to strengthen or weaken the obesity-anxiety relationship.

SEM has been used extensively in public health studies on childhood obesity, typically to identify multilevel risk factors and intervention targets (Forster et al., 2023; Foster et al., 2020). Its relevance to this paper is especially high, since the NSCH data focus on variables representing a variety of ecological levels, such as individual (gender, age, weight status), family (parental education, household income, parental mental health), and community (urbanicity, geographic region). With SEM as the theoretical framework, this study is set to not only add to the body of epidemiological literature in childhood obesity, but also to the body of literature regarding the science of interventions associated with the development of public health, through the identification of ecologically situated high-risk subgroups that have to be targeted through the programming.

Conceptual Framework

Besides the SEM, the study has a conceptual foundation in the emerging literature on psychosocial mechanisms underlying the relationship between childhood obesity and mental health. The existing literature determines that childhood obesity is connected with an increased risk of being exposed to weight-based stigmatization and bullying, lower body esteem, social exclusion, and lower health-related quality of life, all of which are proximal determinants of anxiety and depression (Beltran-Garrayo et al., 2023; Beltran-Garrayo et al.,

2024; Kokka et al., 2023). These psychosocial processes are theoretically cross-located within the ecological levels of the SEM: peer victimization and social comparison are at the microsystem level, attitudes of the family towards weight and mental health of parents are at the mesosystem and exosystem levels, and the cultural norms around the body image and weight gain are at the macrosystem level.

The nature of the conceptual model that guides the present study is that the relationship between childhood obesity, depression, and anxiety is mediated by both gender as bodies image concerns, social comparison processes, and help-seeking behaviors vary by gender and non-gender as access to mental health services, physical activity facilities, and community social support varies systematically in rural and urban settings (Contreras et al., 2021; Foster et al., 2020; Papadimitriou et al., 2025). This theoretical framework aligns with the SEM and allows the development of theoretically significant hypotheses regarding differences in moderation. The research questions will aim to test the hypothesis that these theoretically hypothesized moderation effects are supported by empirical evidence from the U.S. nationally representative NSCH data.

Figure 2 Sociological Model Applied to this study

Nature of the Study

The selected research design for the study will be a quantitative, cross-sectional design based on secondary analysis of publicly available data from the 2022-2023 National Survey of Children's Health (NSCH). This study would be conducted quantitatively, as the research questions will involve statistical testing of associations and the effects of interactions among specifically defined variables, and the hypotheses will be directional and can be tested

using inferential tests. The ⁴ cross-sectional design is suitable because the NSCH ⁹⁰ is used to collect data at a single point in time and is targeted for prevalence and association analyses of a population in the context of pediatric health research (Crouch et al., 2023; Foster et al., 2020). ¹⁵

It is also supported by the fact that the choice of the quantitative paradigm is predetermined by the already existing literature, which has been using epidemiological and biostatistical techniques to study the relationships between obesity ⁹¹ and mental health issues in pediatric patients without exception (Forster et al., 2023; Galler et al., 2024; Kokka et al., 2023; Papadimitriou et al., 2025). ¹⁵ Cross-sectional ⁴ secondary analysis is a type of research that enables the researcher to make robust statistical inferences on a national scale without incurring the costs, ethical issues, and time involved in primary data collection. It ⁴ also allows the use of NSCH survey weights, without which generating nationally representative estimates that can be projected to all children in the target age range in the U.S. is unachievable ⁹²

The main limitation of the cross-sectional design is that it does not consider the cause-and-effect relationships between childhood obesity ⁹³ and anxiety. The ⁴ primary and secondary research questions will be tested using logistic regression with interaction terms. In ⁴ particular, regressions of the independent variable (BMICLASS) on the dependent variables (anxiety and depression) will have interaction terms of gender x BMICLASS and urbanicity x BMICLASS. Each ⁴ model will incorporate the following covariates (depression or anxiety as a cross-control, age, race/ethnicity, socioeconomic ⁷ status) and will be adjusted using NSCH survey weights to make them nationally representative. ⁹⁴

Literature Search Strategy

The literature search strategy that was used in this study was thorough. Accessed ⁴ databases included PubMed, PsycINFO, CINAHL, ERIC, Web of

Science, ProQuest, and Google Scholar, which were used to access the main databases. Walden University Library was used as the primary source for peer-reviewed academic articles. The search terms and combinations were: childhood obesity, pediatric obesity, BMI classification, anxiety disorders in children, depression in children, mental health and obesity, gender differences in obesity-related mental health, urbanicity and child health, National Survey of Children's Health (NSCH), moderation analysis, logistic regression and survey-weighted analysis. Terms were systematically combined using Boolean operators (AND, OR). To be current and relevant, the literature search was restricted to peer-reviewed empirical research published since 2019. Theoretical works, including Bronfenbrenner's (1979) original writings on the social-ecological model, were incorporated regardless of publication date, given their fundamental role in the theory. Systematic reviews and meta-analyses prioritized synthesis of evidence on important associations. The use of grey literature, dissertations, and conference proceedings was sparse, and only when the peer-reviewed material did not cover a particular aspect of the research questions.

Theoretical Framework

The theoretical framework for this research is Bronfenbrenner's Social-Ecological Model (SEM). This framework was first proposed by Bronfenbrenner (1979) in his famous book, Ecology of Human Development, which postulated that human development and behavior result from the dynamic interaction between the individual and various levels of the surrounding world. The main theoretical hypotheses of the SEM are since extensively generalized into the field of public health research as an organizing framework to understand the multilevel determinants of health behaviors and outcomes; (a) individuals do not merely receive health and behavioral outcomes but are the active causal

agents of their ecological context; (b) proximal environments (microsystems like family and school) interact with more distal contexts (exosystems and macrosystems like community resources, cultural norms, and policy frameworks); and (c) health and behavioral outcomes are not merely implied by In this context, the gender is an individual level factor, which modifies how children perceive and react to weight-related stigma, whereas urbanicity is a structural factor at the community level that defines the availability of physical activity, healthy food, and access to mental health care. The moderators (gender and urbanicity) are factors at the individual level (individual) and community level (community), respectively, and covariates (age, race/ethnicity, socioeconomic status) are factors that reflect further ecological interactions. This theoretical stance informs the inclusion of interaction terms in the logistic regression models to test the hypothesis that there are ecological subgroups that differ in the degree of interaction between obesity and anxiety. SEM has been utilized in numerous previous studies on childhood obesity and health outcomes. Using the NSCH data to test the hypothesis that parental mental health was a differentiating factor for child obesity across rural and urban settings and income status, Foster et al. (2020) utilized a multilevel analysis framework based on SEM concepts. Similarly, Forster et al. (2023) used a socio-ecological approach to explain the relationships between BMI and mental health across age groups, genders, and socioeconomic statuses in a German pediatric cohort. The use of the NSCH by Crouch et al. (2023) also allowed the authors to demonstrate that rural-urban differences in childhood obesity persist even after controlling for demographic covariates, which aligns with the SEM assumption that community-level variables have an independent effect on health outcomes. To start, the multilevel SEM design fits the study design, which focuses on individual (gender) and community (urbanicity)

moderators simultaneously. Second, the NSCH dataset includes variables operationalizing various ecological levels, making it a perfect empirical platform for testing the SEM-derived hypotheses. Third, the SEM has a history of research on childhood obesity, providing theoretical continuity with the current body of evidence.

Conceptual Framework

The conceptual basis of the current study is extensively supported by the available research that has recorded the nature of psychosocial processes through which childhood obesity predisposes a greater risk of anxiety and depression. The principle of this framework is weight-based stigmatization and its consequences, i.e., lower body esteem, social exclusion, victimization by peers, and lower quality of life (related to health). These psychosocial stressors play the role of proximal determinants of mental health outcomes. They are theorized to be more intense and to have greater impact with respect to the child's gender and geographic location. Primary writings by major researchers define the conceptual background. The researchers Beltran-Garrayo et al. (2023) proved the conceptual pathway obesity-psychosocial stigmas-anxiety, as the body esteem of the patients and the levels of peer teasing mediated the significantly high levels of anxiety in the obese youth. Beltran-Garrayo et al. (2024) extended this model to a longitudinal level, demonstrating that body esteem mediates the obesity-depression relationship among adolescents. Kokka et al. (2023) have offered a system synthesis that psychiatric comorbidities, such as anxiety, are disproportionately high among obese children working across different study designs. Galler et al. (2024) also established that anxiety disorders are one of the most prevalent psychiatric comorbidities in clinically obese youth and that these comorbidities have a negative impact on treatment outcomes. The following are the conceptual

propositions of the framework: (a) childhood obesity¹¹⁰ leads to higher exposure to weight-related stigma and social comparison processes; (b) these psychosocial stressors have a negative effect on the treatment outcomes¹¹¹; (c) the intensity of these actions depends on gender (boys and girls are different in the way they internalize body image messages. These⁴ propositions are consistent with and generalized to the SEM by defining the psychosocial mechanisms according to which¹¹² ecological variables are converted into¹¹² individual mental health outcomes.

The use of this conceptual framework in previous studies has been a similar^{113,114} manner.¹¹⁴ Papadimitriou et al. (2025)⁴ established that the association between obesity¹¹⁵ and anxiety was mediated by gender and rural-urban dwelling among the young children of a European sample. In⁴ a cohort of nearly 700,000 Germans, Forster et al. (2023)²⁴ identified a gender and socioeconomic⁷ status-dependent mental health relationship with BMI.

Literature Review Related to Key Variables and/or Concepts

The current literature consistently supports a relationship between childhood obesity¹¹⁶ and mental health outcomes, particularly anxiety, while also highlighting important contextual factors. Several⁴ studies using longitudinal and cross-sectional designs (e.g., Beltrán-Garrayo et al., 2023, 2024; Förster et al., 2023)¹⁵ demonstrate that higher BMI is associated with increased psychological distress, often mediated by body esteem and social experiences such as teasing. These⁴ findings justify the selection of obesity¹¹⁷ as the independent variable and anxiety/depression as dependent variables.¹¹⁸ Additionally⁴, studies using NSCH data (Crouch et al., 2023; Foster et al., 2020)¹⁵ validate the appropriateness of a quantitative, cross-sectional methodology with large-scale survey data.

Covariates such as gender, socioeconomic status, and parental mental health are supported by prior findings indicating their influence on both obesity and mental health outcomes. For instance, Papadimitriou et al. (2025) and Contreras et al. (2021) highlight differences across gender and urbanicity, supporting their inclusion as moderators. While most studies report a positive association between obesity and mental health disorders, some inconsistencies remain, particularly regarding anxiety (Kokka et al., 2023), indicating ongoing debate.

Overall, the literature confirms significant associations but reveals a gap in simultaneously examining gender and urbanicity within a single model. This study extends prior research by integrating these moderators to better address the proposed research questions.

Definitions

All major variables considered in this research are operationalized with reference to the 2022–2023 NSCH data dictionary and codebook (Health Resources and Services Administration & U.S. Census Bureau, 2024).

Childhood Obesity (Independent Variable): A BMI at the 95th percentile at age and sex, according to CDC growth charts (Centers for Disease Control and Prevention, 2022). In the NSCH, this is obtained through the variable BMICLASS that classifies children as healthy, overweight and obese with reference to parent-reported height and weight (Health Resources and Services Administration & U.S. Census Bureau, 2024). In this study, BMICLASS will be categorized as obese (BMI ≥95th percentile)

Anxiety (Primary Dependent Variable): The operationalization of anxiety can be explained as a parent-reported symptom with current anxiety disorder, which is based on the NSCH survey item where the parent is asked whether a doctor or health professional has told the parent his/her child has anxiety, and whether

the child has the condition now (Health Resources and Services Administration & U.S. Census Bureau, 2024). This^{4,122} is a dichotomous (yes/no) variable.

Depression (Secondary Dependent Variable): This is analogous in operationalization to anxiety; it is based on the value of the current diagnosis of depression, as reported by parents on the NSCH (Health Resources and Services Administration & U.S. Census Bureau, 2024). It⁴ is also a dichotomous (yes/no) variable.

Gender (Moderator Variable): Measured as a dichotomous variable using the NSCH item on which respondents are asked to report on the sex of the child (male/female), and is congruent with the measurement method of the survey (Health Resources and Services Administration & U.S. Census Bureau, 2024).

Urbanicity (Moderator Variable): The geographical location of the child based on the NSCH metropolitan statistical area (MSA) vs. non-MSA (rural) in line with the rural-urban classification in the same analyses of previous studies with NSCH data (Crouch et al., 2023; Foster et al., 2020)¹⁵.

Socioeconomic Status⁷ (Covariate): Operationalized as a percentage of the federal poverty level (FPL) of household income, a continuous variable in the NSCH, with four levels of analysis (less than 100% FPL, 100–199% FPL, 200–399% FPL and 400% or more FPL) (Health Resources and Services Administration & U.S. Census Bureau, 2024).

Race/Ethnicity (Covariate): Measured using the NSCH race and ethnicity variables, Non-Hispanic White, Non-Hispanic Black, Hispanic, Non-Hispanic Asian, and Non-Hispanic Other/Multiracial (Health Resources and Services Administration & U.S. Census Bureau, 2024).

Age (Covariate): This is the age of the child in years when the survey is filled¹²³, but in this case, it is limited to children between the ages of 10 and 17 years^{123 123 123 123}.

Depression/Anxiety as Cross-Control Covariate: When anxiety is the outcome variable, depression status is added as a covariate, and vice versa, to adjust for the high co-occurring conditions and isolate the independent relationship of each of these conditions with obesity¹²⁴ (Health Resources and Services Administration & U.S. Census Bureau, 2024).

Assumptions

The focus of the present study is determined by using a country-wide cross-sectional secondary data (NSCH 2022-2023) in investigating the relationship between childhood obesity¹²⁵ and anxiety/depression in U.S. children between the ages of 10 and 17 years, using gender and urbanicity as modifying factors. It⁴ was estimated that the age range of 10-17 was the time of development¹²⁶ when body image concerns, peer social comparison and anxiety disorders are clinically salient and measurable by parent report¹²⁷, and when the psychiatric burden of obesity-related stigma is most acute.

The particular emphasis on gender and urbanicity as two moderators¹²⁸, as opposed to¹²⁸ a wider range of potential moderators, was done to fill the gap in¹²⁸ the literature that had been identified and due to the parsimony of analysis¹²⁸. The reason why the NSCH dataset was chosen is due to the availability of the most comprehensive and rigorously methodologically representative nationally representative data on child health in the U.S., with validated parent-reported mental health diagnoses, measured or reported anthropometric data to classify by BMI, and key demographic variables.¹²⁹

The cross-sectional design limits external validity¹³⁰ of the study because it does not permit drawing¹³⁰ causal conclusions, and the use of secondary data limits the researcher to collecting the variables¹³⁰ that were not measured in the NSCH. The scope of the population will be non-institutionalized, civilian U.S. children aged 10-17 years old¹³¹ with no missing data on significant variables¹³¹. Results⁴ are

not to be extended to younger children (under 10), adults and other populations outside the United States. ⁴ Other ¹³² theoretical models, like the cognitive behavioral model of anxiety ¹³² or the biopsychosocial model, were also taken into consideration ¹³², but not taken as the dominant theoretical framework ¹³², because the SEM provides the best correspondence to multilevel moderators ¹³² of interest. ⁷³

Scope and Delimitations

This research has a number of ¹³³ possible contributions to the ¹³³ existing knowledge in the sphere of the epidemiology ¹³³ of public health and pediatrics. To ⁴ begin with, it builds up on the existing literature of the obesity-anxiety relationship by simultaneously testing the moderating impact of gender and urbanicity on one another using a nationally representative sample in the U.S. Previous research has typically studied these moderators independently or in study sample that is not based in the U.S., which presents a gap in the epidemiological literature that this study fills directly (Papadimitriou et al., 2025). ¹⁵ Second, ⁴ survey-weighted logistic regression with interaction terms is a methodologically sound way of moderation testing ¹³⁴, which could be used as the template in future ¹³⁴ epidemiological studies involving complex survey data. ¹³⁴ ¹³⁴ ¹³⁴ ¹³⁴ ¹³⁴

For practice and policy, the findings of the study ¹³⁵ will be useful as actionable evidence to distinguish the sub-groups ¹³⁵ of children (e.g., rural girls and urban boys). ⁴ This fact is of direct importance in relation to health services provided in schools, pediatric primary care providers and community-based organizations involved in obesity prevention and mental health promotion. ¹³⁶ Among ⁴ the policy implications, there is the possibility to support focused funding distributions ¹³⁷ both in ¹³⁷ rural mental health infrastructure and pediatric health programming that is gender-sensitive. ¹³⁷

This study can be useful ¹³⁸ on several ecological levels ¹³⁸ in order to positively change the situation. Individually⁴, the high-risk subgroups can be identified early, and anxious obese children can be intervened in such cases. At⁴ a communal level, the data can inform the formulation of a culturally and geographically specific community health intervention. The⁴ existence of the moderation effect differences¹³⁹, in turn, justifies equity-based public health policy responses to the combination of geographic and gender-based health disparities in childhood, at the policy level^{139 4}. The⁴ research therefore helps in¹⁴⁰ the greater¹⁴⁰ purpose of health research, which is to minimize health disparities and enhance wellbeing among all children irrespective of their places¹⁴⁰ of residence or sex.

Limitations

This research has several limitations associated with the quantitative, cross-sectional research design and secondary NSCH data. First⁴, the internal validity is constrained, as it is impossible to determine causal relations between childhood obesity¹⁴¹ and anxiety; only associations can be investigated. Use⁴ of parent-reported data generates the risk of measurement error, which interferes with construct validity, especially in subjective variables like anxiety and depression. The⁴ country-representative NSCH data increases the external validity, but the results might not apply beyond the U.S. groups or other cultural backgrounds. Potential⁴ methodological weaknesses include omitted-variable bias. Even⁴ though the major covariates, such as socioeconomic⁷ status, gender, and parental mental health, are covered, other factors of interest (e.g., diet quality, physical activity, or genetic predisposition) might not be adequately captured. Obesity⁴ and anxiety are likely to be affected by confounding variables that may affect observed relationships. Parental⁴ responses may be influenced by biases, including social desirability bias and recall bias. To⁴ address the

problems, the research will ¹⁴²utilize statistical controls and multivariable regression ¹⁴²methods to minimize confounding ¹⁴²effects. ⁴Also, it enhances reliability and consistency by using a large, standardized dataset. ⁴Although it is impossible to remove their limitations, attentive choice of variables, powerful analytical techniques, and transparent reporting will contribute to their reduction and enhance the credibility of results.

Significance

The proposed study makes valuable contributions to knowledge in the field of public health by advancing understanding of the links between childhood ¹⁴³obesity and anxiety within a thorough analytical framework. ⁴By incorporating moderating factors such as gender and urbanicity, the study will be grounded in the existing literature and consistent with the Social Ecological Model, which emphasizes ⁷³multi-level determinants of health outcomes. ⁴This study will be of value to the literature by filling gaps in understanding how contextual and demographic factors affect the obesity-mental health relationship. ⁴Practically and policy-wise, the findings can be used to design specific interventions that could help prevent the occurrence of ¹⁴⁴obesity and anxiety in children. ⁴In particular, classifying high-risk subgroups by gender or geographic location can ¹⁴⁵assist healthcare providers, educators, and policymakers ¹⁴⁵in developing more precise prevention programs. ⁴Evidence-based decision-making can be facilitated by insights from nationally representative NSCH data at both the community and national levels. ⁴The study's implications for positive social change are also evident; it will provide the opportunity to identify vulnerable groups at the earliest stage and involve them in combined strategies for physical and mental treatment. ⁴Combined treatment for ¹⁴⁶obesity and anxiety can enhance overall child welfare, reduce long-term health inequalities, and promote healthier developmental processes. ⁴These contributions do not

override the study's scope and focus on the practical implications of enhancing child health outcomes.

Summary and Conclusions

The literature has shown that there is a strong relationship between childhood ¹⁴⁷obesity and poor mental health, specifically, anxiety and depression. ⁴The most important themes are the importance of psychosocial factors, including body esteem and teasing, and demographic factors, including gender, socioeconomic status, and urbanicity. ⁴Empirical ¹⁴⁸research based on big data, such as the NSCH, justifies the application of quantitative techniques to test these relationships. ⁴Nevertheless, the results can be contradictory, especially regarding the power and stability of the positive correlation between anxiety and ¹⁴⁹obesity. ⁴Although there is a lot of information on the overall association between ¹⁵⁰obesity and mental health, there is less on the interaction among multiple moderating factors. ⁴The available literature tends to investigate either gender or urbanicity separately, and there are gaps in models that address both. ⁴The current paper also fills these discrepancies by investigating the relationship between childhood ¹⁵¹obesity and anxiety, and also considering the factors of gender and urbanicity as moderators. ⁴The method extends existing information by offering more detailed insight into the interplay among these variables in a nationally representative sample. ⁴The results will guide future studies and interventions.

Section 2: Research Design and Data Collection

Introduction

In the section, a detailed description of the research design, data source, sampling methods, variable definitions, and data analysis approaches is provided for this quantitative study, which relied on secondary data and employed a cross-sectional analysis. ^{4,152}This is to ensure methodological rigor and

transparency capable of surmounting the challenge of reproducibility and enabling critical analysis of the study findings. ⁴ All the choices made regarding the methods are explained by compliance with the research questions, the theoretical framework, and the specifics of the NSCH data. ⁴ This part will further be subdivided into the following subsections: research design and rationale; setting and sample; instrumentation; data collection procedures; variable definitions; data analysis plan; ethical considerations; and limitations.

Research Design and Rationale

Secondary data analysis is conducted using a quantitative, cross-sectional design in this research. ⁴ The quantitative design is appropriate because the research questions are developed to test a statistical relationship and to assess the moderate impact of operationally measured variables through inferential statistical analysis. ⁴ A cross-sectional design is appropriate because all the data needed are gathered at a single point in time using the NSCH, which is specifically designed to support population-level association research in childhood health. ⁴ ^{153,155} ^{153,155} ¹⁵⁵ The cross-section data is suitable to support the prevalence-based association testing, which constitutes the principal analytic target of the present study, and have been actively employed in ¹⁵⁵ the previous research on childhood ¹⁵⁴ obesity and mental health (Crouch et al., 2023; Foster et al., 2020; Papadimitriou et al., 2025). ¹⁵ ⁴ The 2022-2023 NSCH can be used as a publicly available, high-quality, nationally representative dataset because it is properly designed and used in a secondary data analysis. ⁴ Secondary analysis will not only address the logistical, financial, and ethical constraints of primary data collection it will also provide a far wider sample than would be possible in ^{156,157} a primary data collection. ¹⁵⁷ ⁴ ¹⁵⁹ As a particular source of data, the NSCH has been used in the literature on childhood ¹⁵⁸ obesity and rural-urban health (Crouch et

al., 2023; Foster et al., 2020), ¹⁵ which makes it reasonable to assume it is ¹⁵⁹ relevant to the research questions.

Methodology

The study sample included U.S. children aged 10-17 years who were not in institutions and were sampled in the 2022-2023 National Survey of Children's Health (Data Resource Center for Child and Adolescent Health [DRC], 2024). ⁴ The NSCH sample comprises over 50,000 children nationwide, and an estimated sub-sample of more than 15,000 after applying the study's inclusion ¹⁶⁰ and exclusion criteria. Inclusion ⁴ criteria were: age 10–17 years, non-institutionalized status, and complete data on BMI classification, anxiety diagnosis, and gender. Exclusion ⁴ criteria were: missing or incomplete responses on any study variable and age outside the 10–17 range. The ⁴ initial data set used a complicated ¹⁶¹ sampling method involving probability sampling to select addresses via list-assisted dialing with random digits, ¹⁶¹ ensuring national representativeness. Such ⁴ a methodology indicates a stratified and clustered sample design that improves coverage of the population and its geographic areas. The ⁴ information was obtained from parent- or caregiver-reported surveys conducted by federal agencies, which ensured homogeneous collection practices. This ⁴ study included participants aged 10-17 years with full information on obesity ¹⁶² status, anxiety/depression diagnosis, gender, urbanicity, and the most important covariates. The ⁴ exclusion criteria were the absence of or unrealistic BMI data and age less than 10-17 years, as the data is already anonymous and requires user registration only. The ⁴ federal control and prior peer-reviewed use lend it credibility. Power ⁴ analysis with G*Power (Faul et al., 2007) ¹⁵ indicated that a sample size of 400 was the minimum required; the available sample size was strong in terms of statistical power ¹⁶³ and thus suitable for this study.

Sampling procedures used by original¹⁶⁴ creators of the data set.

The NSCH uses a complex, probability-based sampling design to be nationally representative^{165 4}. In particular, it employs address-based sampling (ABS) with stratified random sampling, which allows covering both landline and non-landline households. Households⁴ are randomly selected from a national address frame, and one child per household is selected. This^{4,166} is a stratified, cluster sampling plan that enhances coverage in areas and demographic classes. Structured⁴ questionnaires were used to collect data from parents or caregivers, administered online or by mail. This⁴ study will have inclusion criteria as children aged 10-17 with full data on BMI, anxiety, depression, gender and urbanicity. The⁴ exclusion criteria involve the implausibility of BMI and key variables. Data⁴ Resource Centre makes the dataset publicly available, and registration is required⁴. It⁴ does not require any special permission as it is de-identified. NSCH⁴ is a federally funded survey with high reliability and validity, and it has been extensively used in peer-reviewed studies, demonstrating high data quality. This⁴ data is perfect because it has a large, nationally representative sample and includes the variables. A⁴ priori power analysis conducted using G*Power (Faul et al., 2007)¹⁵ with $\alpha = .05$, power = .80, and a small-to-medium effect size (OR = 1.5) indicated a minimum required sample of approximately 400 participants. Because⁴ the NSCH dataset contains a substantially larger nationally representative sample, the available sample exceeds the minimum requirement needed to ensure adequate statistical power.

Operationalization

The Health Resources and Services Administration (HRSA) Maternal and Child Health Bureau and the National Centre¹⁶⁷ for Health Statistics (NCHS) of the CDC administered the 2022-2023 NSCH. The⁴ sampling frame was based on all non-

institutionalized children living in the United States who were between the¹⁶⁸ ages of 17 years.¹⁶⁸ The⁴ sampling method used was a list-assisted random-digit-dial telephone sampling technique, supplemented by address-based sampling to increase coverage among households without landline telephones, resulting in a nationally representative sample. The⁴ present study will include the following inclusion criteria: (1) children aged between 10 and 17 years; (2) full data on BMI CLASS (obesity¹⁶⁹ status); (3) full data on parent-reported anxiety and depression diagnosis; (4) full data on gender (sex), urbanicity, and at least 80% complete data on covariates. The⁴ exclusion criteria will be as follows: (1) children between 0-9 and 18 and above years; (2) data on primary independent or dependent variables will be missing; (3) implausible BMI values given by NSCH data quality flags. The⁴ NSCH data is publicly accessible on the Data Resource Centre on Child and Adolescent Health (DRC) at childhealthdata.org. There⁴ is no need for any special permits or institutional agreements to access the public-use dataset, since it is de-identified and released by the federal agency. The⁴ public-use file does not require a formal data use agreement, but the researcher must be registered in the DRC portal to access and download the dataset.

Data Analysis Plan

IBM SPSS Statistics Version 29 was is¹⁷⁰ to analyze the data. The⁴ first step in data cleaning involved verifying missing data, outliers, and implausible BMI records using NSCH data quality flags. The⁴ cases with large amounts of missing data were dropped, and minor missingness was addressed using list wise¹⁷¹ deletion. Survey⁴ weights were applied to make it national. This⁴ study answered two research questions aiming to determine relationships between obesity¹⁷² and anxiety/depression, and the moderators were gender and urbanicity. Hypotheses⁴ were tested using logistic regression analyses, since the outcomes

were binary. ⁴The effect was moderated by including interaction terms (obesity
¹⁷³ ¹⁷⁵ ¹⁷⁵ ¹⁷⁴ ¹⁷⁵ ⁴ x gender, obesity x urbanicity). ⁴The main assumptions were that the
observations were independent, there was no multicollinearity, and the logit
was linear with respect to continuous predictors. ⁴It tested them using variance
inflation factors, correlation matrices, and the Box-Tidwell procedures. ⁴If
assumptions were not met, transformations or large standard errors were to be
considered. ⁴Where appropriate, comparisons were handled using conservative
significance levels. ⁴The introduction of covariates was informed by previous
literature that stated that they confounded the effect. ⁴The interpretation was
done through odds ratios, 95% confidence intervals, and p-values. ⁴ $\alpha = 0.05$ was
set as the statistical significance level. ⁴This type of analysis can be used to
conduct a strong study of association and free-play effects in the dataset.

Threats to Validity

External validity can be compromised by using U.S.-based NSCH data, which
limits the applicability of these results to other nations or cultures. ⁴But in the
United States, it is enhanced by national representativeness (Crouch et al.,
¹⁵ ⁴2023; Foster et al., 2020). ⁴The level of testing reactivity is low because data was
gathered without any reference to this research.

The threats to internal validity ^{176,178}originates from the use of cross-sectional
¹⁷⁸ ¹⁷⁸ ¹⁷⁸design, which ¹⁷⁸restricts the ability to ¹⁷⁸draw causes and introduces a possibility of
reverse causation between anxiety and ¹⁷⁷obesity (Kokka et al., 2023). ¹⁵Probability
sampling helps to reduce selection bias, but residual confounding can still
exist. ⁴Parent-reported measures are also a source of instrumentation threats,
as they can lead to reporting errors (Foster et al., 2020). ¹⁵ ⁴The issues of construct
validity are associated with the fact that single-item measures of anxiety and
depression are used, which may be restrictive in the depth of measurement
(Galler et al., 2024). ¹⁵ ⁴Multicollinearity or model misspecification may also

compromise the statistical conclusion validity; the diagnostic tests were performed to mitigate these threats.

The study addressed these threats by including established variables, adjusting for major confounders, and using appropriate statistical methods (Forster et al., 2023). ²⁴ Where ¹⁵ necessary, ⁴ sensitivity analyses were done. Although ⁴ limitations are inescapable in some respects, the validity and reliability of findings can be improved through a critical approach to methodological design and analytical rigor.

Ethical Procedures

The federal government issues the 2022-2023 NSCH public-use dataset in a de-identified form. The ⁴ public-use file does not contain any personally identifiable information (PII). Not ⁴ all records are publicly published without first being depersonalized (removal of direct identifiers (names, addresses, Social Security numbers, and specific geographic identifiers below the state level) from the records). The ⁴ NCHS and HRSA conducted de-identification before releasing the data, which is in line with the federal privacy protections under the Privacy Act and the Confidential Information Protection and Statistical Efficiency Act (CIPSEA). Since ⁴ the dataset that ¹⁷⁹ is used by the researcher does not contain any PII, there is no possibility of re-identifying individual participants. The ⁴ data storage and data security measures in this study are: downloaded NSCH dataset will be stored on a password-protected computer that is only accessible to the principal researcher, all data analysis files will be encrypted, no raw data file will be shared with unauthorized persons, and a minimum of five years of data will be kept after the study is completed as per Walden University IRB policy before data is deleted. The ⁴ data files in the NSCH are accessible to any registered ¹⁸⁰ user of the DRC, and are not confidential; however, the researcher will treat the downloaded files as confidential research

data in line with good research practice. Two⁴ important ethical issues apply to this study.

To begin with, since the secondary data used in this research are based on a federal government survey, there is no face-to-face interaction with human respondents. Nevertheless⁴, a valuable ethical implication is the sound stewardship of data, which entails child health information. Although⁴ the data are de-identified, the researcher will still have the ethical responsibility to avoid re-identifying individual participants and to use the data in accordance with the purpose of this study. This^{4, 181} shall be addressed through close adherence⁴ to the NSCH data use terms and Walden University IRB protocols. Second⁴, the research investigates the possibly sensitive health issues (obesity¹⁸³, anxiety, and depression) in children, which raises the question of how the results can be applied in a stigmatizing manner. To⁴ reduce the risk of contributing to weight-based or mental health stigmatization, the researcher will ensure that all dissemination of findings is expressed in non-stigmatizing, person-first¹⁸⁴ language and that the findings are presented in¹⁸⁴ a framework of public health equity.¹⁸⁴

Summary

The study adopted a quantitative and cross-sectional design, where secondary data were utilized through the 2022-2023 NSCH to test the association between childhood obesity¹⁸⁵ and anxiety/depression. The⁴ sample of children aged 10-17 used in the study is nationally representative, which provides a strong basis for analyzing the situation. To⁴ achieve consistency and validity, standardized NSCH measures were used to operationalize variables, and a rigorous data analysis plan was implemented, including logistic regression and moderation analyses to examine relationships and interaction effects. The⁴ major covariates were selected based on previous studies, and the statistical

assumptions were thoroughly examined to ensure correct outcomes. The ¹⁸⁶ methodological controls and diagnostic testing were utilized ¹⁸⁶ to overcome ¹⁸⁶ potential threats to validity. The ⁴ use of de-identified data, safe storage procedures, and responsible reporting were considered ethical priorities. All ⁴ ethical considerations were followed in the study, which ensured ¹⁸⁷ confidentiality, reduced ¹⁸⁷ risk, yielded ¹⁸⁷ relevant results that could be generalized to the entire population, and mitigated ¹⁸⁷ limitations. The ⁴ research strategy supports the study's purpose and research questions, providing a clear structure for investigating the relationship between obesity ¹⁸⁸ and mental health. The ⁴ second section presents the results of the statistical analysis, including both descriptive and hypothesis-testing results.

Section 3: Presentation of the Results and Findings

Section 4: Application to Professional Practice and Implications for Social Change

References

Beltrán-Garrayo, L., Larsen, J. K., Eisinga, R., Vink, J. M., Blanco, M., Graell, M., & Sepúlveda, A. R. (2024). Childhood obesity ¹⁸⁹ and adolescent follow-up depressive symptoms: Exploring a moderated mediation model of body esteem and gender. European Child & Adolescent Psychiatry, 33(8), 2859–2869.

<https://doi.org/10.1007/s00787-023-02348-9>

Beltrán-Garrayo, L., Solar, M., Blanco, M., Graell, M., & Sepulveda, A. R. (2023). Examining associations between obesity ¹⁹⁰ and mental health disorders from childhood to adolescence: A case-control prospective study. Psychiatry Research, 326, 115296. <https://doi.org/10.1016/j.psychres.2023.115296>

Bronfenbrenner, U. (1979). The ecology of human development ¹⁹¹: Experiments by nature and design. Harvard ⁴ University Press.

- Centers for Disease Control and Prevention. (2022). CDC growth charts for the United States: Methods and development. U.S. Department of Health and Human Services. <https://www.cdc.gov/growthcharts>
- Contreras, D. A., Martoccio, T. L., Brophy-Herb, H. E., Horodynski, M., Peterson, K. E., Miller, A. L., & Lumeng, J. C. (2021). Rural–urban differences in body mass index and obesity-related behaviors among low-income preschoolers. Journal of Public Health, 43(4), e637–e644. <https://doi.org/10.1093/pubmed/fdaa160>
- Crouch, E., Abshire, D. A., Wirth, M. D., Hung, P., & Benavidez, G. A. (2023). Rural-urban differences in overweight and obesity, physical activity, and food security among children and adolescents. Preventing Chronic Disease, 20, E92. <https://doi.org/10.5888/pcd20.230136>
- Data Resource Center for Child and Adolescent Health. (2024). 2022–2023 National Survey of Children's Health: Flourishing for children and adolescents, age 6–17 years — Nationwide [Data query results]. Child and Adolescent Health Measurement Initiative, Johns Hopkins Bloomberg School of Public Health. <https://www.nschdata.org/browse/survey/results?q=11070&r=1>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behavior Research Methods, 39(2), 175–191. <https://doi.org/10.3758/BF03193146>
- Förster, L. J., Vogel, M., Stein, R., Hilbert, A., Breinker, J. L., Böttcher, M., & Poulain, T. (2023). Mental health in children and adolescents with overweight or obesity. BMC Public Health, 23(1), 135. <https://doi.org/10.1186/s12889-023-15032-z>
- Foster, B. A., Weinstein, K., Mojica, C. M., & Davis, M. M. (2020). Parental mental health is associated with child overweight and obesity, examined within rural

and urban settings, stratified by income. Journal of Rural Health, 36(1), 27–37. <https://doi.org/10.1111/jrh.12395>

Galler, A., Thönnnes, A., Joas, J., Joisten, C., Körner, A., Reinehr, T., Röbl, M., Schauerte, G., Siegfried, W., Weghuber, D., Weihrauch-Blüher, S., Wiegand, S., Holl, R. W., Prinz, N., & APV Initiative. (2024). Clinical characteristics and outcomes of children, adolescents, and young adults with overweight or obesity and mental health disorders. International Journal of Obesity, 48(3), 423–432. <https://doi.org/10.1038/s41366-023-01449-4>

Health Resources and Services Administration & U.S. Census Bureau. (2024). 2022–2023 National Survey of Children's Health: Public use file codebook. Data Resource Center for Child and Adolescent Health. <https://www.nschdata.org>

Kokka, I., Mourikis, I., & Bacopoulou, F. (2023). Psychiatric disorders and obesity in childhood and adolescence—A systematic review of cross-sectional studies. Children, 10(2), 285. <https://doi.org/10.3390/children10020285>

Papadimitriou, K., Mentzelou, M., Papadopoulou, S. K., Antasouras, G., Deligiannidou, G.-E., Alexatou, O., Ntovoli, A., Psara, E., Papadopoulou, V. G., & Giaginis, C. (2025). Childhood obesity and overweight are associated with a higher risk of depression and anxiety: A cross-sectional study in children aged 6–9 years. Life, 15(6), 968. <https://doi.org/10.3390/life15060968>

Ranjani, H., Mehreen, T. S., Pradeepa, R., Anjana, R. M., & Mohan, V. (2021). Epidemiology of childhood overweight and obesity in India: A systematic review. Indian Journal of Endocrinology and Metabolism, 25(3), 192–200. https://doi.org/10.4103/ijem.IJEM_693_20

Appendix A: Title of Appendix

Field Product #1: Policy Brief Memo

Title:

Addressing Childhood Obesity and Anxiety in U.S. Youth: A Dual-Risk Public Health Priority

Introduction

Childhood ²⁰⁰obesity and anxiety are critical and interrelated public health concerns in the United States. ⁴Evidence from national datasets such as the National Survey of Children's Health (NSCH) shows that ²⁰¹obesity is significantly associated with increased risk of anxiety and depression among children aged 10–17.

Scope of the Problem

Approximately 20% of U.S. children are obese, with increasing prevalence of mental health disorders (Crouch et al., 2023; Contreras et al., 2021). ¹⁵Rural ⁴ children and specific gender groups face disproportionate risks due to structural and social inequities (Foster et al., 2020). ¹⁵

Current Approaches

Current interventions often address ²⁰²obesity and mental health separately. ⁴School nutrition and physical activity programs exist but lack integrated mental health screening (Forster et al., 2023). ²⁴ ¹⁵

Proposed Policy

Implement integrated screening programs ²⁰³combining BMI and mental health ²⁰³assessment in pediatric and school settings. ⁴Expand rural mental health services and develop gender-sensitive interventions.

Major Constituencies

Public health agencies, pediatric providers, schools, policymakers, and community organizations.

Conclusion

Integrated and equity-focused policies are essential to reduce the dual burden of ²⁰⁴obesity and anxiety in children.

References

Contreras, Z. A., et al. (2021)

Crouch, E., et al. (2023)

Forster, M., et al. (2023)

Foster, B. A., et al. (2020)

Field Product #2: Community Health Intervention Plan

Integrated Childhood Wellness Program: Addressing Obesity and Mental Health

Problem Definition

Childhood ²⁰⁵obesity is significantly associated with anxiety and depression, particularly among rural and vulnerable populations (Beltran-Garrayo et al., 2023; Kokka et al., 2023).¹⁵

Goal Setting

Reduce ²⁰⁶obesity prevalence

Improve early mental health detection

Increase access to care

Target Population

Children aged 10–17, especially:

Rural populations

Low socioeconomic groups⁷

High-risk gender subgroups

Intervention Strategies

Integrated screening (BMI + mental health)

School-based health education

Counseling services

Community engagement programs

Implementation Plan

Partner with schools and health departments

Train providers

Use mobile clinics in rural areas

Evaluation Plan

Pre/post health outcomes

Participation rates

Reduction in anxiety/depression

References

Beltran-Garrayo, L., et al. (2023)

Kokka, I., et al. (2023)

Crouch, E., et al. (2023)

Foster, B. A., et al. (2020)

DrPH Field Product #3: Visual Representation of Intervention Framework

SEM Level

Intervention Component

Delivery Mode

Responsible Party

Success Indicator

Microsystem

(Individual)

Physical activity sessions & nutrition education

School-based workshops (weekly)

School nurses, PE teachers

10% BMI z-score reduction at 24 months

Microsystem

(Individual)

CBT skills groups (gender-specific)

Small group, bi-weekly

Licensed counselors

15% GAD-7 score reduction

Mesosystem

(Family)

Parent/caregiver education

Monthly workshops & take-home materials

CHWs, school counselors

25% increase in parent knowledge

Exosystem

(School/Community)

School counselor training in integrated screening

Professional ²⁰⁷development workshops

Public health program coordinator

100% of counselors trained

Macrosystem

(Policy)

Community advocacy for rural health equity

Policy briefs, community forums

DrPH candidates, administrators

Policy brief delivered to ²⁰⁸health department

Note. ⁴CHW = Community Health Worker; CBT = Cognitive Behavioral Therapy;

GAD-7 = Generalized Anxiety Disorder-7; SEM = Social Ecological Model

(Bronfenbrenner, 1979). ¹⁵Delivery ⁴modes and indicators are subject to

adaptation based on community needs assessment ²⁰⁹findings.

DrPH Field Product #4: Community Fact Sheet

□ Key Facts About Childhood Obesity in the U.S.

1.	Mph ,	Improper formatting	Correctness
2.	obesity → Obesity	Custom suggestions with replacements	Style guide
3.	ix → is	Misspelled words	Correctness
4.	. Although; . This; . The; . Weighted; . Results; . If; . It; . A; . Another; . Figure; . Data; . In; . Rural; . Studies; . Obese; . As; . These; . Adolescents; . Population-level; . Kokka; . Anxiety; . Likewise; . However; . Thus; . Its; . With; . Cross-sectional; . Each; . Accessed; . Walden;	Text inconsistencies	Correctness
5.	obesity → Obesity	Custom suggestions with replacements	Style guide
6.	obesity → Obesity	Custom suggestions with replacements	Style guide
7.	socioeconomic; socio-economic; Socioeconomic	Text inconsistencies	Correctness
8.	obesity → Obesity	Custom suggestions with replacements	Style guide
9.	obesity → Obesity	Custom suggestions with replacements	Style guide
10.	This	Intricate text	Clarity
11.	This	Intricate text	Clarity
12.	SocialChange → Social Change	Misspelled words	Correctness
13.	obesity → Obesity	Custom suggestions with replacements	Style guide
14.	obesity → Obesity	Custom suggestions with replacements	Style guide

15.	<i>(Contreras et al., 2021; Crouch et al., 2023); (Contreras et al., 2021; Forster et al., 2023); (Crouch et al., 2023; Foster et al., 2020); (Crouch et al., 2023); (Beltran-Garrayo et al., 2023; Galler et al., 2024; Kokka et al., 2023); (Beltran-Garrayo et al., 2024; Forster et al., 2023); (Foster et...</i>	Citation style options	Correctness
16.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
17.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
18.	<i>In particular, there are not many studies that have investigated the moderating effects of gender and urbanicity concomitantly, which are two factors with different, and possibly interactive effects on the obesity-anxiety relationship.</i>	Paragraph can be improved	Clarity
19.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
20.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
21.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
22.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
23.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
24.	<i>Forster; Förster</i>	Text inconsistencies	Correctness
25.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements
26.	obesity → Obesity	Custom suggestions with replacements	Style guide replacements

27.	obesity → Obesity	Custom suggestions with replacements	Style guide
28.	obesity → Obesity	Custom suggestions with replacements	Style guide
29.	obesity → Obesity	Custom suggestions with replacements	Style guide
30.	obesity → Obesity	Custom suggestions with replacements	Style guide
31.	theconnection → the connection	Misspelled words	Correctness
32.	obesity → Obesity	Custom suggestions with replacements	Style guide
33.	obesity → Obesity	Custom suggestions with replacements	Style guide
34.	<i>In a large German cohort (aged 4-18 years), Forster et al. (2023) discovered that the relationships between BMI and mental health outcomes, such as behavioral problems and health-related quality of life, depended on both gender and socioeconomic status, making it important that these moderators be ...</i>	Paragraph can be improved	Clarity
35.	obesity → Obesity	Custom suggestions with replacements	Style guide
36.	obesity → Obesity	Custom suggestions with replacements	Style guide
37.	<i>Likewise, parental mental health has been identified as a significant contextual variable, differentially predicting obesity risk among children across rural and urban environments and income groups (Foster et al., 2020), underscoring the relevance of considering family and environmental context in...</i>	Paragraph can be improved	Clarity

38.	<i>Regardless of this accumulating evidence, there is an identifiable gap in the literature.</i>	Paragraph can be improved	Clarity
39.	obesity → Obesity	Custom suggestions with replacements	Style guide
40.	obesity → Obesity	Custom suggestions with replacements	Style guide
41.	obesity → Obesity	Custom suggestions with replacements	Style guide
42.	obesity → Obesity	Custom suggestions with replacements	Style guide
43.	obesity → Obesity	Custom suggestions with replacements	Style guide
44.	obesity → Obesity	Custom suggestions with replacements	Style guide
45.	obesity → Obesity	Custom suggestions with replacements	Style guide
46.	<i>The NSCH data have been mostly utilized in studies on the prevalence of obesity and rural-urban differences in weight status (Crouch et al., 2023; Foster et al., 2020), without analyzing mental health outcomes as core dependent variables and the moderate effects of gender and urbanicity on each oth...</i>	Paragraph can be improved	Clarity
47.	obesity → Obesity	Custom suggestions with replacements	Style guide

48.	<i>This research is based on the current literature in the area of childhood obesity and mental health, specifically extending two streams of previous research: moderated mediation models (Beltran-Garrayo et al., 2024) and rural-urban comparative studies (Contreras et al., 2021; Crouch et al., 2023).</i>	Paragraph can be improved	Clarity
49.	<i>It develops these frameworks in two important ways - by using a nationally representative U. S. sample and by introducing a dual moderation, that is, two variables simultaneously moderate the relationship under study.</i>	Paragraph can be improved	Clarity
50.	<i>This</i>	Intricate text	Clarity
51.	obesity → Obesity	Custom suggestions with replacements	Style guide
52.	obesity → Obesity	Custom suggestions with replacements	Style guide
53.	study that tests → analysis of	Paragraph can be improved	Clarity
54.	obesity → Obesity	Custom suggestions with replacements	Style guide
55.	obesity → Obesity	Custom suggestions with replacements	Style guide
56.	BMICLASS → BMI CLASS	Misspelled words	Correctness
57.	<i>In particular, the research will focus on identifying whether the connection between childhood obesity (independent variable: BMICLASS, operationalized as BMI 95th percentile per CDC growth charts) and parent-reported anxiety diagnosis (dependent variable — that is, whether the strength or directio...</i>	Paragraph can be improved	Clarity

58.	obesity → Obesity	Custom suggestions with replacements	Style guide
59.	framework,	Incorrect punctuation	Correctness
60.	obesity → Obesity	Custom suggestions with replacements	Style guide
61.	will	Incorrect verb forms	Correctness
62.	intervention → interventions	Incorrect noun number	Correctness
63.	combined,	Incorrect punctuation	Correctness
64.	obesity → Obesity	Custom suggestions with replacements	Style guide
65.	obesity,	Incorrect punctuation	Correctness
66.	an alternative	Determiner use (a/an/the/this, etc.)	Correctness
67.	obesity → Obesity	Custom suggestions with replacements	Style guide
68.	obesity → Obesity	Custom suggestions with replacements	Style guide
69.	obesity → Obesity	Custom suggestions with replacements	Style guide
70.	obesity → Obesity	Custom suggestions with replacements	Style guide
71.	obesity → Obesity	Custom suggestions with replacements	Style guide
72.	obesity → Obesity	Custom suggestions with replacements	Style guide
73.	multi-level; multilevel	Text inconsistencies	Correctness

74.	development → Development	Custom suggestions with replacements	Style guide
75.	that human	Conjunction use	Correctness
76.	development → Development	Custom suggestions with replacements	Style guide
77.	outcome → outcomes	Incorrect noun number	Correctness
78.),	Incorrect punctuation	Correctness
79.	<i>These propositions lead directly to the study's analytic approach, in which gender is treated as an individual-level moderator, urbanicity as a community-level moderator, and how the two interact to strengthen or weaken the obesity-anxiety relationship.</i>	Paragraph can be improved	Clarity
80.	obesity → Obesity	Custom suggestions with replacements	Style guide
81.	<i>Its relevance to this paper is especially high, since the NSCH data focus on variables representing a variety of ecological levels, such as individual (gender, age, weight status), family (parental education, household income, parental mental health), and community (urbanicity, geographic region).</i>	Paragraph can be improved	Clarity
82.	obesity → Obesity	Custom suggestions with replacements	Style guide
83.	development → Development	Custom suggestions with replacements	Style guide
84.	<i>Besides the SEM, the study has a conceptual foundation in the emerging literature on psychosocial mechanisms underlying the relationship between childhood obesity and mental health.</i>	Paragraph can be improved	Clarity

85.	obesity → Obesity	Custom suggestions with replacements	Style guide
86.	obesity → Obesity	Custom suggestions with replacements	Style guide
87.	obesity → Obesity	Custom suggestions with replacements	Style guide
88.	development → Development	Custom suggestions with replacements	Style guide
89.	study.	Closing punctuation	Correctness
90.	is used to collect → collects	Paragraph can be improved	Clarity
91.	obesity → Obesity	Custom suggestions with replacements	Style guide
92.	unachievable.	Closing punctuation	Correctness
93.	obesity → Obesity	Custom suggestions with replacements	Style guide
94.	<i>Each model will incorporate the following covariates (depression or anxiety as a cross-control, age, race/ethnicity, socioeconomic status) and will be adjusted using NSCH survey weights to make them nationally representative.</i>	Paragraph can be improved	Clarity
95.	obesity → Obesity	Custom suggestions with replacements	Style guide
96.	obesity → Obesity	Custom suggestions with replacements	Style guide
97.	obesity → Obesity	Custom suggestions with replacements	Style guide

98.	<i>The use of grey literature, dissertations, and conference proceedings was sparse, and only when the peer-reviewed material did not cover a particular aspect of the research questions.</i>	Paragraph can be improved	Clarity
99.	development → Development	Custom suggestions with replacements	Style guide
100.	obesity → Obesity	Custom suggestions with replacements	Style guide
101.	obesity → Obesity	Custom suggestions with replacements	Style guide
102.	obesity → Obesity	Custom suggestions with replacements	Style guide
103.	<i>Using the NSCH data to test the hypothesis that parental mental health was a differentiating factor for child obesity across rural and urban settings and income status, Foster et al. (2020) utilized a multilevel analysis framework based on SEM concepts.</i>	Paragraph can be improved	Clarity
104.	obesity → Obesity	Custom suggestions with replacements	Style guide
105.	obesity → Obesity	Custom suggestions with replacements	Style guide
106.	obesity → Obesity	Custom suggestions with replacements	Style guide
107.	a greater	Determiner use (a/an/the/this, etc.)	Correctness
108.	<i>They are theorized to be more intense and to have greater impact with respect to the child's gender and geographic location.</i>	Paragraph can be improved	Clarity

109.	have a negative impact on → hurt, harm	Wordy sentences	Clarity
110.	obesity → Obesity	Custom suggestions with replacements	Style guide
111.	have a negative effect on → hurt, harm	Wordy sentences	Clarity
112.	<i>These propositions are consistent with and generalized to the SEM by defining the psychosocial mechanisms according to which ecological variables are converted into individual mental health outcomes.</i>	Paragraph can be improved	Clarity
113.	in a	Wrong or missing prepositions	Correctness
114.	<i>The use of this conceptual framework in previous studies has been a similar manner.</i>	Paragraph can be improved	Clarity
115.	obesity → Obesity	Custom suggestions with replacements	Style guide
116.	obesity → Obesity	Custom suggestions with replacements	Style guide
117.	obesity → Obesity	Custom suggestions with replacements	Style guide
118.	the dependent	Determiner use (a/an/the/this, etc.)	Correctness
119.	obesity → Obesity	Custom suggestions with replacements	Style guide
120.	obesity → Obesity	Custom suggestions with replacements	Style guide

121.	<i>In the NSCH, this is obtained through the variable BMICLASS that classifies children as healthy, overweight and obese with reference to parent-reported height and weight (Health Resources and Services Administration & U.S. Census Bureau, 2024).</i>	Paragraph can be improved	Clarity
122.	<i>This</i>	Intricate text	Clarity
123.	<i>Age (Covariate): This is the age of the child in years when the survey is filled, but in this case, it is limited to children between the ages of 10 and 17 years.</i>	Paragraph can be improved	Clarity
124.	obesity → Obesity	Custom suggestions with replacements	Style guide
125.	obesity → Obesity	Custom suggestions with replacements	Style guide
126.	development → Development	Custom suggestions with replacements	Style guide
127.	report,	Incorrect punctuation	Correctness
128.	<i>The particular emphasis on gender and urbanicity as two moderators, as opposed to a wider range of potential moderators, was done to fill the gap in the literature that had been identified and due to the parsimony of analysis.</i>	Paragraph can be improved	Clarity
129.	<i>The reason why the NSCH dataset was chosen is due to the availability of the most comprehensive and rigorously methodologically representative nationally representative data on child health in the U.S., with validated parent-reported mental health diagnoses, measured or reported anthropometric data...</i>	Paragraph can be improved	Clarity

130.	<i>The cross-sectional design limits external validity of the study because it does not permit drawing causal conclusions, and the use of secondary data limits the researcher to collecting the variables that were not measured in the NSCH.</i>	Paragraph can be improved	Clarity
131.	<i>The scope of the population will be non-institutionalized, civilian U.S. children aged 10-17 years old with no missing data on significant variables.</i>	Paragraph can be improved	Clarity
132.	<i>Other theoretical models, like the cognitive behavioral model of anxiety or the biopsychosocial model, were also taken into consideration, but not taken as the dominant theoretical framework, because the SEM provides the best correspondence to multilevel moderators of interest.</i>	Paragraph can be improved	Clarity
133.	<i>This research has a number of possible contributions to the existing knowledge in the sphere of the epidemiology of public health and pediatrics.</i>	Paragraph can be improved	Clarity
134.	<i>Second, survey-weighted logistic regression with interaction terms is a methodologically sound way of moderation testing, which could be used as the template in future epidemiological studies involving complex survey data.</i>	Paragraph can be improved	Clarity
135.	<i>For practice and policy, the findings of the study will be useful as actionable evidence to distinguish the sub-groups of children (e.g., rural girls and urban boys).</i>	Paragraph can be improved	Clarity
136.	obesity → Obesity	Custom suggestions with replacements	Style guide

137.	<i>Among the policy implications, there is the possibility to support focused funding distributions both in rural mental health infrastructure and pediatric health programming that is gender-sensitive.</i>	Paragraph can be improved	Clarity
138.	<i>This study can be useful on several ecological levels in order to positively change the situation.</i>	Paragraph can be improved	Clarity
139.	<i>The existence of the moderation effect differences, in turn, justifies equity-based public health policy responses to the combination of geographic and gender-based health disparities in childhood, at the policy level.</i>	Paragraph can be improved	Clarity
140.	<i>The research therefore helps in the greater purpose of health research, which is to minimize health disparities and enhance wellbeing among all children irrespective of their places of residence or sex.</i>	Paragraph can be improved	Clarity
141.	obesity → Obesity	Custom suggestions with replacements	Style guide
142.	<i>To address the problems, the research will utilize statistical controls and multivariable regression methods to minimize confounding effects.</i>	Paragraph can be improved	Clarity
143.	obesity → Obesity	Custom suggestions with replacements	Style guide
144.	obesity → Obesity	Custom suggestions with replacements	Style guide
145.	<i>In particular, classifying high-risk subgroups by gender or geographic location can assist healthcare providers, educators, and policymakers in developing more precise prevention programs.</i>	Paragraph can be improved	Clarity

146.	obesity → Obesity	Custom suggestions with replacements	Style guide
147.	obesity → Obesity	Custom suggestions with replacements	Style guide
148.	provides justification for applying	Paragraph can be improved	Clarity
149.	obesity → Obesity	Custom suggestions with replacements	Style guide
150.	obesity → Obesity	Custom suggestions with replacements	Style guide
151.	obesity → Obesity	Custom suggestions with replacements	Style guide
152.	<i>This</i>	Intricate text	Clarity
153.	<i>The cross-section data is suitable to support the prevalence-based association testing, which constitutes the principal analytic target of the present study, and have been actively employed in the previous research on childhood obesity and mental health (Crouch et al., 2023; Foster et al., 2020; Pa...</i>	Ungrammatical sentence	Correctness
154.	obesity → Obesity	Custom suggestions with replacements	Style guide
155.	<i>The cross-section data is suitable to support the prevalence-based association testing, which constitutes the principal analytic target of the present study, and have been actively employed in the previous research on childhood obesity and mental health (Crouch et al., 2023; Foster et al., 2020; Pa...</i>	Paragraph can be improved	Clarity
156.	collectionit → collection, but it	Conjunction use	Correctness

157.	<i>Secondary analysis will not only address the logistical, financial, and ethical constraints of primary data collection it will also provide a far wider sample than would be possible in a primary data collection.</i>	Paragraph can be improved	Clarity
158.	obesity → Obesity	Custom suggestions with replacements	Style guide
159.	<i>As a particular source of data, the NSCH has been used in the literature on childhood obesity and rural-urban health (Crouch et al., 2023; Foster et al., 2020), which makes it reasonable to assume it is relevant to the research questions.</i>	Paragraph can be improved	Clarity
160.	study's inclusion	Improper formatting	Correctness
161.	<i>The initial data set used a complicated sampling method involving probability sampling to select addresses via list-assisted dialing with random digits, ensuring national representativeness.</i>	Paragraph can be improved	Clarity
162.	obesity → Obesity	Custom suggestions with replacements	Style guide
163.	sufficiently powerful	Paragraph can be improved	Clarity
164.	the original	Determiner use (a/an/the/this, etc.)	Correctness
165.	ensure national representativeness	Paragraph can be improved	Clarity
166.	This	Intricate text	Clarity
167.	Centre → Center	Misspelled words	Correctness
168.	<i>The sampling frame was based on all non-institutionalized children living in the United States who were between the ages of 17 years.</i>	Paragraph can be improved	Clarity

169.	obesity → Obesity	Custom suggestions with replacements	Style guide
170.	is → used	Incorrect verb forms	Correctness
171.	list-wise → listwise	Misspelled words	Correctness
172.	obesity → Obesity	Custom suggestions with replacements	Style guide
173.	obesity → Obesity	Custom suggestions with replacements	Style guide
174.	obesity → Obesity	Custom suggestions with replacements	Style guide
175.	<i>The effect was moderated by including interaction terms (obesity x gender, obesity x urbanicity).</i>	Paragraph can be improved	Clarity
176.	<i>The threats to internal validity originates from the use of cross-sectional design, which restricts the ability to draw causes and introduces a possibility of reverse causation between anxiety and obesity (Kokka et al., 2023).</i>	Ungrammatical sentence	Correctness
177.	obesity → Obesity	Custom suggestions with replacements	Style guide
178.	<i>The threats to internal validity originates from the use of cross-sectional design, which restricts the ability to draw causes and introduces a possibility of reverse causation between anxiety and obesity (Kokka et al., 2023).</i>	Paragraph can be improved	Clarity
179.	<i>Since the dataset that is used by the researcher does not contain any PII, there is no possibility of re-identifying individual participants.</i>	Passive voice misuse	Clarity
180.	user of the DRC, → DRC user	Paragraph can be improved	Clarity

181.	<i>This</i>	Intricate text	Clarity
182.	<i>by adhering closely</i>	Paragraph can be improved	Clarity
183.	obesity → Obesity	Custom suggestions with replacements	Style guide
184.	<i>To reduce the risk of contributing to weight-based or mental health stigmatization, the researcher will ensure that all dissemination of findings is expressed in non-stigmatizing, person-first language and that the findings are presented in a framework of public health equity.</i>	Paragraph can be improved	Clarity
185.	obesity → Obesity	Custom suggestions with replacements	Style guide
186.	<i>The methodological controls and diagnostic testing were utilized to overcome potential threats to validity.</i>	Paragraph can be improved	Clarity
187.	<i>All ethical considerations were followed in the study, which ensured confidentiality, reduced risk, yielded relevant results that could be generalized to the entire population, and mitigated limitations.</i>	Paragraph can be improved	Clarity
188.	obesity → Obesity	Custom suggestions with replacements	Style guide
189.	obesity → Obesity	Custom suggestions with replacements	Style guide
190.	obesity → Obesity	Custom suggestions with replacements	Style guide
191.	development → Development	Custom suggestions with replacements	Style guide
192.	development → Development	Custom suggestions with replacements	Style guide

193.	obesity → Obesity	Custom suggestions with replacements	Style guide
194.	obesity → Obesity	Custom suggestions with replacements	Style guide
195.	obesity → Obesity	Custom suggestions with replacements	Style guide
196.	obesity → Obesity	Custom suggestions with replacements	Style guide
197.	obesity → Obesity	Custom suggestions with replacements	Style guide
198.	obesity → Obesity	Custom suggestions with replacements	Style guide
199.	obesity → Obesity	Custom suggestions with replacements	Style guide
200.	obesity → Obesity	Custom suggestions with replacements	Style guide
201.	obesity → Obesity	Custom suggestions with replacements	Style guide
202.	obesity → Obesity	Custom suggestions with replacements	Style guide
203.	<i>Implement integrated screening programs combining BMI and mental health assessment in pediatric and school settings.</i>	Paragraph can be improved	Clarity
204.	obesity → Obesity	Custom suggestions with replacements	Style guide
205.	obesity → Obesity	Custom suggestions with replacements	Style guide

206.	obesity → Obesity	Custom suggestions with replacements	Style guide
207.	development → Development	Custom suggestions with replacements	Style guide
208.	the health	Determiner use (a/an/the/this, etc.)	Correctness
209.	<i>Delivery modes and indicators are subject to adaptation based on community needs assessment findings.</i>	Paragraph can be improved	Clarity